Network Systems
Science & Advanced
Computing

Biocomplexity Institute & Initiative

University of Virginia

Estimation of COVID-19 Impact in Virginia

May 11th, 2022

(data current to May 7th – 10th)
Biocomplexity Institute Technical report: TR BI-2022-1121



BIOCOMPLEXITY INSTITUTE

biocomplexity.virginia.edu

About Us

- Biocomplexity Institute at the University of Virginia
 - Using big data and simulations to understand massively interactive systems and solve societal problems
- Over 20 years of crafting and analyzing infectious disease models
 - Pandemic response for Influenza, Ebola, Zika, and others



Points of Contact

Bryan Lewis brylew@virginia.edu

Srini Venkatramanan srini@virginia.edu

Madhav Marathe marathe@virginia.edu

Chris Barrett@virginia.edu

Model Development, Outbreak Analytics, and Delivery Team

Przemyslaw Porebski, Joseph Outten, Brian Klahn, Alex Telionis,
Srinivasan Venkatramanan, Bryan Lewis,
Aniruddha Adiga, Hannah Baek, Chris Barrett, Jiangzhuo Chen, Patrick Corbett,
Stephen Eubank, Galen Harrison, Ben Hurt, Dustin Machi, Achla Marathe,
Madhav Marathe, Mark Orr, Akhil Peddireddy, Erin Raymond, James Schlitt, Anil Vullikanti,
Lijing Wang, James Walke, Andrew Warren, Amanda Wilson, Dawen Xie



Overview

• Goal: Understand impact of COVID-19 mitigations in Virginia

Approach:

- Calibrate explanatory mechanistic model to observed cases
- Project based on scenarios for next 4 months
- Consider a range of possible mitigation effects in "what-if" scenarios

Outcomes:

- Ill, Confirmed, Hospitalized, ICU, Ventilated, Death
- Geographic spread over time, case counts, healthcare burdens

Key Takeaways

Projecting future cases precisely is impossible and unnecessary. Even without perfect projections, we can confidently draw conclusions:

- Case rates continue to rise as pace picks up as do hospitalizations
- VA 7-day mean daily case rate increased to 26/100K from 18.5/100K
 - US continues to increase slightly to 22/100K (from 18.5/100K)
 - VA hospital occupancy (rolling 7 day mean of 240) has steadily rising for nearly a month
- Surveillance anomalies continue as QA processes rebalance previously reported cases though is slowing
- Projections anticipate future growth in cases but more limited growth in more severe outcomes:
 - Recently emerging BA.2.12.1 subvariant seems to driving growth, VA has tracked BA.2.12.1 scenario of April 23rd projections closely
 - Potential for significant number of infections remains high, uncertainty surrounds impact of weather and changing social interactions

Model updates:


- Adjusted fitting to work on district level to reduce biases from limited outbreaks within counties and surveillance anomalies
- Adaptive scenario BA.2.12.1 scenario to capture the future growth of this more transmissible variant
- Models need to change their focused outcome to hospitalization or aggregate counties to districts to minimize noisy fluctuations

The situation continues to change. Models continue to be updated regularly.

Situation Assessment



Case Rates (per 100k) and Test Positivity




District Trajectories

Goal: Define epochs of a Health District's COVID-19 incidence to characterize the current trajectory

Method: Find recent peak and use hockey stick fit to find inflection point afterwards, then use this period's slope to define the trajectory

Hockey stick fit




Trajectory	Description	Weekly Case Rate (per 100K) bounds
Declining	Sustained decreases following a recent peak	below -0.9
Plateau	Steady level with minimal trend up or down	above -0.9 and below 0.5
Slow Growth	Sustained growth not rapid enough to be considered a Surge	above 0.5 and below 2.5
In Surge	Currently experiencing sustained rapid and significant growth	2.5 or greater




District Trajectories – last 10 weeks

Status	# Districts (prev week)
Declining	2 (5)
Plateau	0 (2)
Slow Growth	11 (17)
In Surge	22 (11)

Curve shows smoothed case rate (per 100K) Trajectories of states in label & chart box Case Rate curve colored by Reproductive number






MUNIVERSITY of VIRGINIA

CDC's new COVID-19 Community Levels

What Prevention Steps Should You Take Based on Your COVID-19 Community Level?

Low	Medium	High	
 Stay <u>up to date</u> with COVID-19 vaccines <u>Get tested</u> if you have symptoms 	 If you are at high risk for severe illness, talk to your healthcare provider about whether you need to wear a mask and take other precautions Stay up to date with COVID-19 vaccines Get tested if you have symptoms 	 Wear a mask indoors in public Stay up to date with COVID-19 vaccines Get tested if you have symptoms Additional precautions may be needed for people at high risk for severe illness 	
People may choose to mask at any time. People with symptoms, a positive test, or exposure to someone with COVID-19 should wear a mask.			

COVID-19 Community Levels – Use the Highest Level that Applies to Your Community				
New COVID-19 Cases Per 100,000 people in the past 7 days	Indicators	Low	Medium	High
	New COVID-19 admissions per 100,000 population (7-day total)	<10.0	10.0-19.9	≥20.0
Fewer than 200	Percent of staffed inpatient beds occupied by COVID-19 patients (7-day average)	<10.0%	10.0-14.9%	≥15.0%
	New COVID-19 admissions per 100,000 population (7-day total)	NA	<10.0	≥10.0
200 or more	Percent of staffed inpatient beds occupied by COVID-19 patients (7-day average)	NA	<10.0%	≥10.0%





The COVID-19 community level is determined by the higher of the new admissions and inpatient beds metrics, based on the current level of new cases per 100,000 population in the past 7 days



CDC Data Tracker Portal

CDC's new COVID-19 Community Levels





Red outline indicates county had 200 or more cases per 100k in last week


Pale color indicates either beds or occupancy set the level for this county

Dark color indicates both beds and occupancy set the level for this county

COVID-19 Community Levels – Use the Highest Level that Applies to Your Community				
New COVID-19 Cases Per 100,000 people in the past 7 days	Indicators	Low	Medium	High
	New COVID-19 admissions per 100,000 population (7-day total)	<10.0	10.0-19.9	≥20.0
Fewer than 200	Percent of staffed inpatient beds occupied by COVID-19 patients (7-day average)	<10.0%	10.0-14.9%	≥15.0%
	New COVID-19 admissions per 100,000 population (7-day total)	NA	<10.0	≥10.0
200 or more	Percent of staffed inpatient beds occupied by COVID-19 patients (7-day average)	NA	<10.0%	≥10.0%

The COVID-19 community level is determined by the higher of the new admissions and inpatient beds metrics, based on the current level of new cases per 100,000 population in the past 7 days

District Trajectories with Community Levels





Estimating Daily Reproductive Number –

Redistributed gap

May 9th Estimates

Region	Date Confirmed R _e	Date Confirmed Diff Last Week
State-wide	1.081	0.057
Central	1.116	0.096
Eastern	1.026	-0.083
Far SW	1.009	0.177
Near SW	1.040	0.049
Northern	1.090	0.039
Northwest	1.107	0.268






Skipping Weekend Reports & holidays biases estimates
Redistributed "big" report day to fill in gaps, and then estimate R from
"smoothed" time series

Methodology

- Wallinga-Teunis method (EpiEstim¹) for cases by confirmation date
- Serial interval: updated to discrete distribution from observations (mean=4.3, Flaxman et al, Nature 2020)
- Using Confirmation date since due to increasingly unstable estimates from onset date due to backfill




^{1.} Anne Cori, Neil M. Ferguson, Christophe Fraser, Simon Cauchemez. A New Framework and Software to Estimate Time-Varying Reproduction Numbers During Epidemics. American Journal of Epidemiology, Volume 178, Issue 9, 1 November 2013, Pages 1505–1512, https://doi.org/10.1093/aje/kwt133

Mask Usage and Vaccination

Self-reported mask usage continues to fall

- US and VA experienced similar decreases
- Vaccination has leveled off and has leveled off after a slight rise in early April






Data Source: https://covidcast.cmu.edu

Wastewater Monitoring

Wastewater provides a coarse early warning of COVID-19 levels in communities


- Overall in the US, there is an increase in sites with increased levels of virus compared to 15 days ago, however the pace of growth slows
- Current virus levels are at or exceeding max of previous historical levels, has slowed, though more sites are entering upper quintiles





COVID-like Illness Activity

COVID-like Illness (CLI) gives a measure of COVID transmission in the community

- Emergency Dept (ED) based CLI is more correlated with case reporting
- Urgent Care (UC) is more sensitive and is a leading indicator but is prone to some false positives
- As testing behaviors and case ascertainment levels shift, these measures may capture disease better than confirmed cases
- Current trends in UC CLI are slightly up







SARS-CoV2 Variants of Concern

Emerging new variants will alter the future trajectories of pandemic and have implications for future control


- Emerging variants can:
 - Increase transmissibility
 - Increase severity (more hospitalizations and/or deaths)
 - Limit immunity provided by prior infection and vaccinations
- Genomic surveillance remains very limited
 - Challenges ability to estimate impact in US to date and estimation of arrival and potential impact in future


WHO label	Pango lineage•	GISAID clade	Nextstrain clade	Additional amino acid changes monitored°	Earliest documented samples	Date of designation
Alpha	B.1.1.7	GRY	20I (V1)	+S:484K +S:452R	United Kingdom, Sep-2020	18-Dec-2020
Beta	B.1.351	GH/501Y.V2	20H (V2)	+S:L18F	South Africa, May-2020	18-Dec-2020
Gamma	P.1	GR/501Y.V3	20J (V3)	+S:681H	Brazil, Nov-2020	11-Jan-2021
Delta	B.1.617.2	G/478K.V1	21A, 21I, 21J	+S:417N +S:484K	India, Oct-2020	VOI: 4-Apr-2021 VOC: 11-May-2021
Omicron*	B.1.1.529	GRA	21K, 21L	+R346K	Multiple countries, Nov-2021	VUM: 24-Nov-2021 VOC: 26-Nov-2021

Omicron Prevalences subvariant BA.2 dominates

CDC nowcast for week ending May 7th shows 99% overall BA.2 in Region 3 with BA 2.12.1 at 48%

Overall BA.2 in USA now at 99% (BA.2.12.1 at 43%)










SARS-CoV2 Omicron and Sub-Variants

As detected in whole Genomes in public repositories



VoC Polynomial Fit Projections







Note: Data lags force projections to start in past. Everything from dotted line forward is a projection.


Influence of Previous Waves on Next Wave

Variability in timing and intensity of some variant-driven waves influence the timing and severity of subsequent waves

- Some regions had early Delta waves, others lagged; subvariants of Omicron have similar variability
- Not all states are in sync within their HHS regions
- Note outliers for Omicron subvariants








18


Influence of Previous Waves on Next Wave

Strong Delta wave leads to delayed Omicron BA.1 wave

• At US state level (right), Region 6 (AK, TX, OK, LA, NM) was hit early and hard with Delta, however, Omicron arrived later than in Region 2 (NY, NJ) which had a negligible Delta wave



Proportion of cases since July by state



Similar pattern plays out in the regions of VA


- The Southwest had a strong Delta wave and a more delayed Omicron BA.1 wave
- The Northern region had very minimal cases during Delta but experienced an early and strong Omicron BA.1 wave
- Current uptick in Northern regions from BA.2 and BA.2.12.1 may presage a delayed wave in Southwest as well

Pandemic Pubs

- **1.** Observed escape of BA.4 and BA.5 from BA.1 immunity is not as severe as BA.1 against previous immunity but low absolute neutralization levels for BA.4 and BA.5, particularly in the unvaccinated group, indicate low protection against infection
- **2.** BA.4 and BA.5 have weaker ACE2 binding than BA.2 which may result in lower intrinsic transmissibility
- **3.** Estimated US seroprevalence increased from 33.5% to 57.7% from December 2021 to February 2022; higher among children
- **4.** A systematic review of Long-Covid studies show prevalence from 51-80% for mild to severe infections


Pooled mean prevalence results for any experience of PACS, extracted from nine systematic reviews, ranged from 51%–80%.


-Evidence indicates PACS is a condition experienced by a substantial number of individuals with previous SARS-CoV-2 infection. Care for patients with PACS will likely place added stresses on health care and social support systems, including increased emergency department visits, outpatient care, inpatient care and rehabilitation involving multidisciplinary teams.



Researchers in SA isolated live BA.4 and BA.5 viruses and tested them against neutralizing immunity elicited to BA.1 infection in participants who were Omicron/BA.1 infected but unvaccinated (n=24) and participants vaccinated with Pfizer BNT162b2 or Johnson and Johnson Ad26.CoV.2S with breakthrough Omicron/BA.1 infection (n=15)

ttps://www.medrxiv.org/content/10.1101/2022.04.29.22274477v





Researchers show variants BA.2.12.1, BA.4 and BA.5 have increased immune evasion relative to BA.2 but some associated mutations result in weaker ACE2 binding which may lead to reduced intrinsic transmissibility.

https://www.researchsquare.com/article/rs-1611421/v1

https://www.publichealthontario.ca/-/media/Documents/nCoV/ipac/2022/04/post-acute-covid-syndrome-pacs.pdf?sc_lang=en

United States Case Rates

 Rebounding activity, mainly in the Northeast


States

8 (14)

22 (21)

6 (9)

18 (10)



Slow Growth

Status


Declining

Plateau

In Surge

Virginia and Her Neighbors


- All have dramatically dropped from peaks
- Rates have moderated
- All but Kentucky are below 10/100K



United States Hospitalizations


- Hospital admissions are lagging case rates, and have mainly entered plateaus
- Rebounds in the Northeast seen with some rising hospitalization rates

Status	# States
Declining	1 (7)
Plateau	21 (26)
Slow Growth	23 (13)
In Surge	5 (4)




County-level comparison to previous highest peak

Recent Incidence Compared to Worst Week by County




Recent Incidence Compared to Worst Week by County




County-level comparison to last Summer

- Most counties in VA have had the highest case rate of the pandemic in the last week
- Nationally the number of counties at their highest rate has expanded considerably

Recent Incidence Compared to Weekly Summer Mean by County Mean: 12.45; Median: 0.72; IQR: 0.21-2.25








Zip code level weekly Case Rate (per 100K)

Case Rates in the last week by zip code

 Some counts are low and suppressed to protect anonymity, those are shown in white




Based on Spatial Empirical Bayes smoothed point prevalence for week ending 2022-05-09.



Risk of Exposure by Group Size and HCW prevalence

Case Prevalence in the last week by zip code used to calculate risk of encountering someone infected in a gathering of randomly selected people (group size 25)


- **Group Size**: Assumes 2 undetected infections per confirmed case (ascertainment rate from recent seroprevalence survey), and shows minimum size of a group with a 50% chance an individual is infected by zip code (eg in a group of 55 in Fort Eustis, there is a 50% chance someone will be infected)
- **HCW ratio**: Case rate among health care workers (HCW) in the last week using patient facing health care workers as the denominator / general population's case prevalence



Current Hot-Spots

Case rates that are significantly different from neighboring areas or model projections


- **Spatial**: Getis-Ord Gi* based hot spots compare clusters of zip codes with weekly case prevalence higher than nearby zip codes to identify larger areas with statistically significant deviations
- **Temporal**: The weekly case rate (per 100K) projected last week compared to observed by county, which highlights temporal fluctuations that differ from the model's projections



Scenario Trajectory Tracking

Which scenario from last projection did each county track closest?

- Minimal difference between projections overall
- State level trend tracking BA.2.12.1 scenarios (red and green), but not all, likely due to variation in prevalence across the state






Model Update – Adaptive Fitting



Model Structure Extended for Multiple Strains

Omicron escapes immunity from vaccinated and those infected with Delta

- Multiple strain support allows representation of differential protection based on immunological history
- Severity of outcomes varies by strain and level of immunity, thus allowing model to better capture hospitalizations and deaths from Omicron
- Adaptive fitting approach continues to use simulation to generate the full distribution of immune states across the population



Adaptive Fitting Approach

Each county fit precisely, with recent trends used for future projection


 Allows history to be precisely captured, and used to guide bounds on projections


Model: An alternative use of the same meta-population model, PatchSim with multiple tiers of immunity

- Allows for future "what-if" Scenarios to be layered on top of calibrated model
- Allows for waning of immunity and for partial immunity against different outcomes (eg lower protection for infection than death)

External Seeding: Steady low-level importation

 Widespread pandemic eliminates sensitivity to initial conditions, we use steady 1 case per 10M population per day external seeding







Using Ensemble Model to Guide Projections

Ensemble methodology that combines the Adaptive with machine learning and statistical models such as:


- Autoregressive (AR, ARIMA)
- Neural networks (LSTM)
- Kalman filtering (EnKF)


Weekly forecasts done at county level.


Models chosen because of their track record in disease forecasting and to increase diversity and robustness.


Ensemble forecast provides additional 'surveillance' for making scenario-based projections.


Also submitted to CDC Forecast Hub.











Seroprevalence updates to model design


Several seroprevalence studies provide better picture of how many actual infections have occurred

 CDC Nationwide Commercial Laboratory Seroprevalence Survey

Pre-Omicron these findings were consistent with an ascertainment ratio of ~2-3x

- Thus there were 2.5 total infections in the population for every confirmed case recently
- Case ascertainment for Omicron infections are half of that for pre-Omicron, thus for every case there are ~5 total infections
- During the peak of Omicron, the degradation of test seeking and capacity were modeled to have fallen by 3x with a rebound to pre-Omicron levels by mid-Feb

New Data released late on April 26th not yet incorporated in model



Virginia

Feb 22nd: 45% [42% - 48%]; Jan 22nd: 34% [31%-39%]





https://covid.cdc.gov/covid-data-tracker/#national-lab

Calibration Approach

- Data:
 - County level case counts by date of onset (from VDH)
 - Confirmed cases for model fitting
- **Calibration:** fit model to observed data and ensemble's forecast
 - Tune transmissibility across ranges of:
 - Duration of incubation (5-9 days), infectiousness (3-7 days)
 - Undocumented case rate (1x to 7x) guided by seroprevalence studies
 - Detection delay: exposure to confirmation (4-12 days)
 - Approach captures uncertainty, but allows model to precisely track the full trajectory of the outbreak
- **Project:** future cases and outcomes generated using the collection of fit models run into the future
 - Mean trend from last 7 days of observed cases and first week of ensemble's forecast used
 - Outliers removed based on variances in the previous 3 weeks
 - 2 week interpolation to smooth transitions in rapidly changing trajectories
- **Outcomes**: Data driven by shift and ratio that has least error in last month of observations
 - Hospitalizations: 3 days from confirmation, 6.8% of cases hospitalized
 - Deaths: 11 days from confirmation, 1.45% of cases die





Confirmed[†]

COVID-19 in Virginia:





Dashboard Updated: 5/10/2022 Data entered by 5:00 PM the prior day.

		Cases, Hospitaliza	ntions and Deaths		
otal C	ases*	Total H Admiss		Tot Dea	
0	,, ,,	50,4	404	20,295	
w Cases	s: 2,484)^	,		,	
1†	Probable†	Confirmed†	Probable†	Confirmed†	Probable†

* Includes both people with a positive test (Confirmed), and symptomatic with a known exposure to COVID-19 (Probable).

^New cases represent the number of confirmed and probable cases reported to VDH in the past 24 hours.

† VDH adopted the updated CDC COVID-19 confirmed and probable surveillance case definitions on August 27, 2020. Found

here: https://wwwn.cdc.gov/nndss/conditions/coronavirus-disease-2019-covid-19/case-definition/2020/08/05/

Source: Cases - Virginia Electronic Disease Surveillance System (VEDSS), data entered by 5:00 PM the prior day

Outl	breaks
Total Outbreaks* 7,778	Outbreak Associated Cases
	131,054

* At least two (2) lab confirmed cases are required to classify an outbreak

487,643

Testing (PCR Only)		
Testing Encounters PCR Only*	Current 7-Day Positivity Rate PCR Only**	
13,579,268	12.9%	

* PCR" refers to "Reverse transcriptase polymerase chain reaction laboratory testing."

^{**} Lab reports may not have been received yet. Percent positivity is not calculated for days with incomplete data.

Multisystem Inflammatory Syndrome in Children	
Total Cases*	Total Deaths
176	1

*Cases defined by CDC HAN case definition: https://emergency.cdc.gov/han/2020/han00432.asp

Accessed 8:15am May 11, 2022

https://www.vdh.virginia.gov/coronavirus/

^{**} Hospitalization of a case is captured at the time VDH performs case investigation. This underrepresents the total number of hospitalizations in

Scenarios – Transmission Conditions


- Variety of factors continue to drive transmission rates
 - Seasonal impact of weather patterns, travel and gatherings, fatigue and premature relaxation of infection control practices
- Waning Immunity: Mean of 6 months to a year protection (rate of 0.0027) similar to <u>Pfizer study</u>, Omicron waning with a mean of 4 months
- Projection Scenarios:
 - Adaptive: Control remains as is currently experienced into the future with assumption that Omicron remains as the majority strain, and that infection with Omicron provides protection against Omicron infection in the future
 - Adaptive-VariantBA2_12: Same as Adaptive, but with BA.2.12.1 subvariant
 continuing predominance and having a 30% transmission advantage over existing
 Omicron (mainly the overall BA.2 subvariant)
 - Adaptive-VariantBA2_12-IncreasedControl: Same as Adaptive-VariantBA2_12, but with a 25% reduction in transmission to increased mitigations starting in 30 days and phasing into full effect over 2 weeks




Scenarios – Omicron BA.2.12 Description

BA.2.12.1 shows signs of increased transmissibility via increasing prevalence in the US, especially the Northeast

- Transmissibility: Not as well observed as previous VoCs as mainly in US and worldwide genotyping efforts have slowed
- Using a 30% boost to transmissibility
- **Prevalence**: Growth rate compared to BA.2 seems to be similar as to BA.2's vs. BA.1 (and BA.1.1), thus assuming similar prevalence curve (30% growth advantage, doubling ~every 8 days)
- Conservatively estimating prevalence to hit 50% on June 1st with ~95% 4 weeks later
- **Severity:** Assumed to be same as for other Omicron subvariants



Coarse Polynomial fitted estimates of BA.2.12.1 prevalence



Projection Scenarios – Combined Conditions

Name	Txm Controls	Vax	Description
Adaptive	С	SQ	Likely trajectory based on conditions remaining similar to the current experience, includes immune escape due to Omicron
Adaptive-VariantBA2	С	SQ	Transmission rates for BA.2 infections are 30% more infectious, BA.2 prevalence reached 50% on April 1st and rises to over 95% by mid-May
Adaptive-VariantBA2_12	С	SQ	Transmission rates for BA.2.12.1 infections are and additional 30% higher, with BA.2.12.1 prevalence reaching 50% on June 1 st and rising to ~95% 4 weeks after
Adaptive-VariantBA2_12-IncreasedControl	Increased	SQ	Same as Adaptive-VariantBA2_12 with increased mitigations reducing transmission by 25% starting June 1st

Transmission Controls: C = Current levels persist into the future

Increased = Transmission rates are reduced by 25% over 2 weeks starting May 1st

Spring = Transmission rates from mid-Jan 2021 through mid-March 2021 are coarsely replayed,

representing a 60% reduction in transmission rate drivers, with Omicron remaining dominant


Vaccinations: SQ = Status quo acceptance leads to low rates of vaccination through the summer

VO = Vaccination acceptance optimistically expands with increased rates through the summer

13-May-22 38

Last projection comparison – two weeks ago

- Growth to present still tracking April 23rd BA.2.12.1 scenario (orange dashed)
- Solid lines are current projections, dashed lines are from 2 weeks ago
- Current projections not significantly different from previous projection




Model Results




Outcome Projections


Confirmed cases



Estimated Hospital Occupancy




Daily Deaths



Death ground truth from VDH "Event Date" data, most recent dates are not complete

Daily Hospitalized




^{*} without surveillance correction VariantBA2 peaked over 10K in July



Outcome Projections – Closer Look


Confirmed cases




* without surveillance correction VariantBA2 peaked over 10K in July



Daily Hospitalized




Daily Deaths




Death ground truth from VDH "Event Date" data, most recent dates are not complete

Detailed Projections: All Scenarios


Projections by Region



Projections by District



Daily confirmed cases) by rate (per 100K) District (grey with 7-day average in black) with simulation colored by scenario




Alleghany


Eastern Shore

Detailed Projections: All Scenarios - Closer Look


Projections by Region



Projections by District



Daily confirmed cases by rate (per 100K) District (grey with 7-day average in black) with simulation colored by scenario




Hospital Demand and Bed Capacity by Region


Capacities by Region

COVID-19 capacity ranges from 80% (dots) to 120% (dash) of total beds

Adaptive



Adaptive – Variant BA2 12



Length of Stay more variable with Omicron, occupancy projections may vary as a result,

Estimated LOS shortened to better fit observed data

Projections show continued declines and with expanded capacities and adjusted length of stay, no capacities exceeded

ad-hoc estimation performed per region

Interactive Dashboard with regional projections https://nssac.bii.virginia.edu/covid-19/vmrddash/



Length of Stay Estimates

Central

Eastern

Far SW

Near SW


Northern


Northwestern

Current Influenza Hospitalization Forecast

Statistical models for submitting to CDC FluSight forecasting challenge


Hospitalizations nationwide are rising, VA still steady





CDC FluSight **Ensemble Forecasts**

Hospital Admissions for Influenza and Forecast for next 4 weeks (UVA ensemble)





Key Takeaways

Projecting future cases precisely is impossible and unnecessary. Even without perfect projections, we can confidently draw conclusions:

- Case rates continue to rise as pace picks up as do hospitalizations
- VA 7-day mean daily case rate increased to 26/100K from 18.5/100K
 - US continues to increase slightly to 22/100K (from 18.5/100K)
 - VA hospital occupancy (rolling 7 day mean of 240) has steadily rising for nearly a month
- Surveillance anomalies continue as QA processes rebalance previously reported cases though is slowing
- Projections anticipate future growth in cases but more limited growth in more severe outcomes:
 - Recently emerging BA.2.12.1 subvariant seems to driving growth, VA has tracked BA.2.12.1 scenario of April 23rd projections closely
 - Potential for significant number of infections remains high, uncertainty surrounds impact of weather and changing social interactions

Model updates:

- Adjusted fitting to work on district level to reduce biases from limited outbreaks within counties and surveillance anomalies
- Adaptive scenario BA.2.12.1 scenario to capture the future growth of this more transmissible variant
- Models need to change their focused outcome to hospitalization or aggregate counties to districts to minimize noisy fluctuations

The situation continues to change. Models continue to be updated regularly.

Additional Analyses



Overview of relevant on-going studies

Other projects coordinated with CDC and VDH:


- Scenario Modeling Hub: Consortium of academic teams coordinated via MIDAS / CDC to that provides regular national projections based on timely scenarios
- **Genomic Surveillance:** Analyses of genomic sequencing data, VA surveillance data, and collaboration with VA DCLS to identify sample sizes needed to detect and track outbreaks driven by introduction of new variants etc.
- Mobility Data driven Outreach locations: Collaboration with VDH state and local,
 Stanford, and SafeGraph to leverage anonymized cell data to help identify sites most frequently visited by different demographic groups


COVID-19 Scenario Modeling Hub – Round 13

Collaboration of multiple academic teams to provide national and state-by-state level projections for 4 aligned scenarios

- Round 13 results getting finalized
 - Scenarios: New Variant in Summer and waning compared (yes/no new variant vs. 4 month or 10 month waning)
- Prelim results shared internally
- Only national consortium tracking
 Omicron wave well
- Rounds 4-12 now available Round 4 Results were published
 May 5th, 2021 in MMWR

https://covid19scenariomodelinghub.org/viz.html





13-May-22 50

Busiest Places: Mobility Data Can Assist

SafeGraph provides fine-grained mobility measures


- <u>SafeGraph</u>: anonymized geolocation data aggregated from numerous cell phone apps
- One of the most fine-grained and high-coverage mobility data sources available: 6.4 million POIs in the US; 158,869 POIs in VA
- Has been utilized by hundreds of researchers, governments, and the CDC to aid COVID-19 efforts (Chang, Pierson, Koh, et al., <u>Nature 2020</u>; Chang et al, KDD 2021)
- Daily and hourly number of visits to points-of-interest (POIs), i.e., nonresidential locations such as restaurants, bars, gas stations, malls, grocery stores, churches, etc.
- Weekly reports per POI of *where visitors are coming from* (at the census block group level)
- Still has <u>limitations</u> to be aware of (e.g., less representation among children and seniors)






Find the Busiest Locations

POIs are individual addresses, need some aggregation to busy areas






Busiest locations vary by day of week (and time of day)

Find locations visited by Target Population

Census Block Groups in Danville



- Use census data to characterize the populations of the different census block groups
- Identify most frequently visited POIs for each CBG
- 3. Cluster most visited POIs
- 4. Provide potential sites grouped by the demographic groups they likely serve

Goal: Provide frequently visited locations based on populations and vaccination levels one desires to reach

Example: List of locations in the Southside frequented by Black Virginians

Overview of the current roster of targeted populations

These are the current roster of targeted population groups that we are providing as part of the weekly delivery to VDH. (This roster is subject to change.)

- Whole population (eg, no target population filters are applied)
- Race Black
- Ethnicity Latinx
- Ages 20-40
- Ages 20-30
- Ages 30-40
- Unvaccinated populations
- Latinx or Black

Data Elements in the CSV

Rank & LocationWeight
The LocationWeight is estimated #
of visits to POIs in the L14 from the
target group. Rank indicates the
order from most- to 25th mostvisited

HighlyVisitedAddress
This is the address of the POI in the L14 that sees the most visits. It is provided to make it easier to find the L14 on the map.

AreaMostVisitedPeriod
This is the 4-hour period in
the week when the L14 sees
its highest traffic. This is not
target group-specific

Population Group For a targeted file like this one, these will all be the same value.

> S2 Key (L14)

AreaMostVisitedDay
This is the day of the week
when most visitors go to this
S2 location. This is not target
group-specific.

Lat and Lon
This is the latitude
and longitude for
the center of the
L14.

	$\overline{}$		n	÷۱	,
C	U	u	ш	T)	/

VDH District

Locality	District	PopulationGroup	LocationID	Rank Lo	cationWeight	AreaMostVisitedDay	HighlyVisitedAddress	AreaMostVisitedPeriod	Lat	Lon
Accomack Co	Eastern Shor	Latinx or Black	89ba2b55	1	4966.030095	Friday	25297 Lankford Hwy Rt 13 N, (Friday 17:00-21:00	37.6978738	-75.716796
Accomack Co	Eastern Shor	Latinx or Black	89ba2caf	2	3728.476605	Friday	26036 Lankford Hwy, Onley, VA	Friday 15:00-19:00	37.6881681	-75.722612
Accomack Co	Eastern Shor	Latinx or Black	89ba2b57	3	3508.193676	Saturday	25274 Lankford Hwy, Onley, VA	Saturday 13:00-17:00	37.69859	-75.722612
Accomack Co	Eastern Shor	Latinx or Black	89bbd4ad	4	2582.802769	Wednesday	25102 Lankford Hwy, Onley, VA	Sunday 11:00-15:00	37.7023677	-75.710981
Accomack Co	Eastern Shor	Latinx or Black	89ba2b53	5	1844.868961	Sunday	25102 Lankford Hwy, Onley, VA	Friday 16:00-20:00	37.7030842	-75.716796
Albemarle C	Blue Ridge	Latinx or Black	89b38647	1	14088.0684	Thursday	1215 Lee St, University of Virg	Thursday 07:00-11:00	38.0327733	-78.500766
Albemarle C	Blue Ridge	Latinx or Black	89b477ff	2	6999.363545	Saturday	1980 Rio Hill Ctr, Charlottesvill	Saturday 12:00-16:00	38.087391	-78.472353
Albemarle C	Blue Ridge	Latinx or Black	89b38645	3	5824.383454	Wednesday	Cabell Hall 525 Mccormick Roa	Wednesday 11:00-15:00	38.033334	-78.506447
Albemarle C	Blue Ridge	Latinx or Black	89b3888d	4	5078.488029	Friday	540 Pantops Ctr, Pantops, VA,	Thursday 11:00-15:00	38.0334982	-78.455301
Albemarle C	Blue Ridge	Latinx or Black	89b387fd	5	4655.844131	Saturday	100 Twentyninth Place Ct, Cha	Saturday 11:00-15:00	38.077516	-78.478036

Mobility Data Updated Weekly

Box: https://virginia.box.com/s/03kq8el0kzd9w43wz2g3myozov76uizo


Excel sheets and simple HTML maps packaged for use

VDH has a dashboard available upon request to allow interactive viewing

https://arcgis.vdh.virginia.gov/portal/apps/opsdashboard/index.html#/8

631cfc4f181460fafc7e1923f41d581

 Dashboard is restricted to VDH offices and those who VPN into the CoV Network



References

Venkatramanan, S., et al. "Optimizing spatial allocation of seasonal influenza vaccine under temporal constraints." *PLoS Computational Biology* 15.9 (2019): e1007111.

Arindam Fadikar, Dave Higdon, Jiangzhuo Chen, Bryan Lewis, Srinivasan Venkatramanan, and Madhav Marathe. Calibrating a stochastic, agent-based model using quantile-based emulation. SIAM/ASA Journal on Uncertainty Quantification, 6(4):1685–1706, 2018.

Adiga, Aniruddha, Srinivasan Venkatramanan, Akhil Peddireddy, et al. "Evaluating the impact of international airline suspensions on COVID-19 direct importation risk." *medRxiv* (2020)

NSSAC. PatchSim: Code for simulating the metapopulation SEIR model. https://github.com/NSSAC/PatchSim

Virginia Department of Health. COVID-19 in Virginia. http://www.vdh.virginia.gov/coronavirus/

Biocomplexity Institute. COVID-19 Surveillance Dashboard. https://nssac.bii.virginia.edu/covid-19/dashboard/

Google. COVID-19 community mobility reports. https://www.google.com/covid19/mobility/

Biocomplexity page for data and other resources related to COVID-19: https://covid19.biocomplexity.virginia.edu/



Questions?

Points of Contact

Bryan Lewis brylew@virginia.edu

Srini Venkatramanan srini@virginia.edu

Madhav Marathe marathe@virginia.edu

Chris Barrett@virginia.edu

Biocomplexity COVID-19 Response Team

Aniruddha Adiga, Abhijin Adiga, Hannah Baek, Chris Barrett, Golda Barrow, Richard Beckman, Parantapa Bhattacharya, Jiangzhuo Chen, Clark Cucinell, Patrick Corbett, Allan Dickerman, Stephen Eubank, Stefan Hoops, Ben Hurt, Ron Kenyon, Brian Klahn, Bryan Lewis, Dustin Machi, Chunhong Mao, Achla Marathe, Madhav Marathe, Henning Mortveit, Mark Orr, Joseph Outten, Akhil Peddireddy, Przemyslaw Porebski, Erin Raymond, Jose Bayoan Santiago Calderon, James Schlitt, Samarth Swarup, Alex Telionis, Srinivasan Venkatramanan, Anil Vullikanti, James Walke, Andrew Warren, Amanda Wilson, Dawen Xie

